

Baryons, Boundaries and Matrix Models.

Iosif Bena¹, Radu Roiban² and Radu Tatar³¹ Department of Physics
University of California
Los Angeles, CA 90095
iosif@physics.ucla.edu² Department of Physics
University of California
Santa Barbara, CA 93106
radu@vulcan2.physics.ucsb.edu³ Department of Physics
and
Theoretical Physics Group
Lawrence Berkeley National Laboratory
University of California
Berkeley, CA 94720
rtatar@socrates.berkeley.edu**Abstract**

A natural extension of the Dijkgraaf-Vafa proposal is to include fields in the fundamental representation of the gauge group. In this paper we use field theory techniques to analyze gauge theories whose tree level superpotential is a generic polynomial in multi-trace operators constructed out of such fields. We show that the effective superpotential is generated by planar diagrams with at most one (generalized) boundary. This justifies the proposal put forward in [12].

We then proceed to extend the gauge theory-matrix model duality to include baryonic operators. We obtain the full moduli space of vacua for an $U(N)$ theory with N flavors. We also outline a program leading to a string theory justification of the gauge theory-matrix model correspondence with fundamental matter.

1 Introduction

Recently, Dijkgraaf and Vafa have proposed [1, 2, 3] a perturbative method for computing the effective glueball superpotential of several classes of $N = 1$ theories. The proposal instructs one to compute the planar free energy of the matrix model whose potential is the tree-level superpotential of the theory of interest.

Being obtained via a “string theory route”, the original proposal naturally includes theories with fields transforming in the adjoint/bifundamental representation of the gauge group. One of the most natural extensions of this duality is to theories with fields in the fundamental representation of the gauge group. It turns out that this goal can be reached in a rather simple way: one needs to compute only the contribution to the free energy of the matrix model arising from Feynman diagrams with one boundary [9, 12]. Thus, the gauge theory effective superpotential is constructed as

$$W_{eff}(S, \Lambda) = N_c S \left(1 - \ln \frac{S}{\Lambda^3}\right) + N_c \frac{\partial F_{\chi=2}}{\partial S} + N_f F_{\chi=1} \quad . \quad (1)$$

Such a construction was successfully used to compare matrix model predictions with known gauge theory results for theories with massive and massless flavors, with $\mathcal{N} = 1$ and $\mathcal{N} = 2$ supersymmetry [9, 12, 11, 13, 17, 27]. Other interesting related work has appeared in [10, 14, 28, 26, 18, 19, 20, 21, 23, 24, 22].

Of a much deeper interest has been the investigation of the structure underlying this duality. In [4] it was proven that in a gauge theory with adjoint fields, the computation of the superpotential is reproduced by the matrix computation, diagram by diagram. Using anomaly-based arguments, in [15] it was also shown that the gauge theory and the matrix model results are related as proposed by Dijkgraaf and Vafa.

In this paper we prove using field theory techniques that the effective superpotential of a theory with a tree level superpotential being a generic polynomial in multi-trace and baryonic operators is generated by planar diagrams with at most one (generalized) boundary. This justifies the proposal put forward in [12]. We then proceed to extend the gauge theory-matrix model duality to include baryonic operators ¹. We obtain the full moduli

¹As we were preparing the manuscript, [25] appeared which has some overlap with our discussion of baryons.

space of vacua for an $SU(N)$ theory with N flavors. We then outline a program for the string theory justification of the extensions of the Dijkgraaf-Vafa duality.

2 Field theory analysis of the effective superpotential

In this section we analyze in detail the effective superpotential as a function of the glueball superfield $S = \text{Tr}(W^2)$ and the various coupling constants that exist in the theory, and show that in theories with fields transforming in the fundamental representation of the gauge group the superpotential is generated entirely by Feynman diagrams with a single boundary. The analysis is similar to the one described in [15]. We will first consider tree level superpotentials built out of traces of products of quark bilinears. This analysis proves the proposal put forward in [12], that only diagrams with one boundary contribute to the effective superpotential. We will then add multi-trace deformations as well as baryonic operators.

We begin by considering a theory with a fairly arbitrary, polynomial, tree level superpotential:

$$W_G = \sum_{l \geq 1} g_l \text{Tr}[(Q\tilde{Q})^l] \quad , \quad (2)$$

where the trace is over the flavor indices. If no tree level superpotential is present, the global symmetry of the theory is $SU(N_f) \times SU(N_f) \times \prod_{i=1}^{N_f} U(1)_i \times \prod_{i=1}^{N_f} \tilde{U}(1)_i \times U(1)_R$. The introduction of the superpotential above, with scalar couplings, breaks this symmetry group to

$$SU(N_f) \times U(1)_i \times \tilde{U}(1)_i \times U(1)_R \quad . \quad (3)$$

As in the case of a theory with adjoint fields [15], these symmetries are not sufficient to completely determine the effective superpotential. This is easy to see by considering the following rather unusual charge assignment for the various fields and couplings:

	$SU(N_f)$	$U(1)$	$\widetilde{U}(1)$	$U(1)_R$	Δ	
$Q :$	N_f	1	0	1	1	
$\tilde{Q} :$	1	0	1	1	1	
$g_l :$	1	$-l$	$-l$	$2 - 2l$	$3 - 2l$	
$S :$	1	0	0	2	3	(4)

The last column in the table above represents the engineering dimension of the corresponding field.

A simple counting argument shows that all higher loop² Feynman diagrams with insertions of the glueball superfield are finite. We are thus looking for a function which depends only on S and g_i . Furthermore, since this function is generated perturbatively, neither one of its arguments is allowed to appear at a fractional power. It is then not hard to see that the basic invariant combinations are

$$\frac{g_l S^{l-1}}{g_1^l} \quad (\forall) \quad l \geq 2 \quad (5)$$

and they also have vanishing scaling dimension. Thus, the most general effective superpotential which can be generated is

$$W_{eff} = S F\left(\frac{g_l S^{l-1}}{g_1^l}\right) \quad (6)$$

where F is an arbitrary analytic function of its arguments³. A generic term in the series expansion of W_{eff} is

$$c_{l_1 \dots l_n} \left(\prod_{i=1}^n \frac{g_{l_i}}{g_1^{l_i}} \right) S^{1 + \sum_i (l_i - 1)} \quad , \quad (7)$$

where the l_i $i = 1, \dots, n$ are not necessarily distinct and $c_{l_1 \dots l_n}$ are numerical coefficients. It is clear from the coupling constant dependence that this term is generated by a Feynman diagram with n vertices with the corresponding couplings g_{l_i} $i = 1, \dots, n$. The S -dependence of this term also fixes the number of index loops in the diagram. Indeed, since S involves a trace, it is clear that each index loop will generate at most one factor of the glueball superfield. This implies that the diagram will have at least

$$L = 1 + \sum_{i=1}^n (l_i - 1) \quad (8)$$

²The one loop graphs are the only divergent ones; their contribution can be taken into account using anomaly arguments [15]. An attempt in this direction was made in [26]

³The appearance of inverse powers of g_1 is allowed perturbatively, as it can be accommodate by rescaling the quark fields. Indeed, absorbing $\sqrt{g_1}$ in both Q and $\bar{Q}$ implies that the relevant couplings to consider are $\frac{g_l}{g_1}$.

index loops. As in the case of diagrams with fields transforming in the adjoint representation of the gauge group, it is possible to show that, for a Feynman diagram with vertices generated by (2), which has genus g and b boundaries, the number of gauge index loops is given by

$$L_g = 2 - 2g - b + \sum_{i=1}^n (l_i - 1) \quad . \quad (9)$$

Thus, since the theory we are interested in has no fields in the adjoint representation (and thus each diagram has at least one boundary), the only way for the relevant Feynman diagrams to have the number of index loops required by (8) is to be planar and have exactly one boundary. This proves the proposal put forward in [12].

It is easy to add multi-trace deformations to the tree level superpotential in equation (2). The coupling constants now carry several indices, counting the number of traces as well as the number of quark bilinears in each trace. Thus, the terms with l quark bilinears are:

$$\sum_{s_1+\dots+s_m=l} g_{s_1\dots s_m}^{(l)} \text{Tr}[(Q\tilde{Q})^{s_1}] \dots \text{Tr}[(Q\tilde{Q})^{s_m}] \quad (10)$$

It is clear that the charges of $g_{s_1\dots s_m}^{(l)}$ under the symmetry group (3) are the same as the charges of g_l in the table above. Thus, the generalization of (5) to this case is

$$\frac{g_{s_1\dots s_m}^{(l)} S^{l-1}}{g_1^l} \quad (\forall) \quad l \geq 2, \quad (\forall) \quad m \quad . \quad (11)$$

As in the previous case, the Feynman diagrams generating the term proportional to $S^{1+\sum_i(l_i-1)}$ must have $1 + \sum_i(l_i - 1)$ index gauge loops. Unfortunately, the relation between the type of vertices used in the diagram, the number of gauge index loops and the number of boundaries becomes more complicated than (9). It is however not hard to see that the number of index loops remains unchanged if one replaces a single-trace vertex with a multiple-trace vertex while keeping the genus unchanged.

Graphically, this corresponds to a degeneration of a disk by pinching its boundary, as is shown in figure 1. On the left-hand-side of that figure we show a vacuum diagram constructed out of a 4-point and a 6-point single-trace vertices. The outer line represents the boundary (corresponding to a trace over flavor indices) while the inner lines represent gauge index loops. On the right-hand-side we show two possible degenerations of this disk. Following

route (a), we replace the single-trace 4-point vertex by a double-trace one. Following route (b), we replace the single-trace 6-point vertex by a double-trace one. Since these surfaces lie at the boundary of the moduli space of




Figure 1: Degenerations of the disk.

the disk with holes, it is clear from this perspective that the latter diagrams should contribute on the same footing as the former ones. With this definition of boundary components, it follows that equation (9) still applies and the diagrams contributing to the superpotential have exactly one boundary.

Let us now turn to including baryonic operators

$$B = \det Q \quad \tilde{B} = \det \tilde{Q} \quad (12)$$

if $N_f = N_c$ and

$$\begin{aligned} B^i &= \epsilon_{\alpha_1 \dots \alpha_{N_c}} Q_{i_1}^{\alpha_1} \dots Q_{i_{N_c}}^{\alpha_{N_c}} \epsilon^{i_1 \dots i_{N_c} \dots i} \\ \tilde{B}_i &= \epsilon^{\alpha_1 \dots \alpha_{N_c}} \tilde{Q}_{\alpha_1}^{i_1} \dots \tilde{Q}_{\alpha_{N_c}}^{i_{N_c}} \epsilon_{i_1 \dots i_{N_c} \dots i} \end{aligned} \quad (13)$$

if $N_f = N_c + 1$. We will repeat the analysis in the beginning of this section and construct the basic invariants which can be built out of these operators. We will deform the initial theory only by terms linear in the baryons. More general deformations can be treated similarly. The charges of the baryon sources are then:

	$SU(N_f)$	$U(1)$	$\widetilde{U(1)}$	$U(1)_R$	Δ	
$b :$	1	$-N_c$	0	$2 - N_c$	$3 - N_c$	(14)
$\tilde{b} :$	1	0	$-N_c$	$2 - N_c$	$3 - N_c$	

It is easy to see that the basic invariants are

$$I_{l,n} = \frac{g_l}{g_1^l} \left[\frac{b}{g_1^{N_c/2}} \frac{\tilde{b}}{g_1^{N_c/2}} \right]^n S^{(l-1)+n(N_c-2)} \quad (15)$$

where n is an arbitrary number. Thus, the superpotential is a series expansion in $I_{l,n}$. We can then immediately see the Feynman diagram origin of a generic term in this series:

$$S \left[\prod_i \frac{g_i}{g_1} S^{l-1} \right] \left[\frac{b}{g_1^{N_c/2}} \frac{\tilde{b}}{g_1^{N_c/2}} S^{N_c-2} \right]^{n_i} . \quad (16)$$

If all n_i vanish, this term matches equation (7) and each color index loop contributes one power of S . Consider then adding a baryon and anti-baryon to the diagram giving this contribution. To produce a 1PI diagram we need to break one of the initial color index loops and connect it to one Q and one $\tilde{Q}$ from the baryons. Thus, we find that the number of index loops increases with $N_c - 2$ if the quark lines are connected in a planar way. Adding more baryon-anti-baryon pairs follows the same pattern and recovers the second factor in equation (16).

We would like to point out that the notion of planarity we use here is slightly more general than usual. Due to the presence of the Levi-Civita symbols at the baryon vertices one is allowed to use their symmetry properties to reorganize the internal lines of a diagram.

With hindsight, this result is not surprising in light of the discussion concerning multi-trace deformations of the initial theory. Since baryon-anti-baryon pairs can be re-expressed as sums of multi-trace operators, it follows that the counting of powers of S for correlation functions involving insertions of such pairs is identical to the counting of powers of S for correlation functions involving insertions of multi-trace operators. There we saw that each color index loop gave a single S and the diagram was required to be planar and have a single boundary in the generalized sense described above.

We have thus proven that in theories with a tree-level superpotential containing multi-trace operators and/or baryon operators, the effective superpotential as a function of the glueball superfield is generated by planar Feynman diagrams with a single boundary in the generalized sense defined above.

3 On the dynamics of massless fields

One of the most puzzling aspects of the Dijkgraaf-Vafa proposal relating a gauge theory with a matrix model is the fact that nonperturbative physics in the former can be obtained by doing perturbative computations in the latter.

To relate the two theories, we are instructed to first compute the free energy of the matrix model and construct from it the gauge theory effective superpotential. As it is the case with all dualities, there exists a dictionary between objects on both sides, but these objects are not the same. One can only compare correlation functions involving these objects, but not the objects themselves. Therefore, to make contact between the two theories, one needs to integrate out all fields and to compare the effective superpotentials as (complicated) functions of the coefficients of the tree level superpotential and the dynamically generated scale.

It seems therefore that the only way one can compute gauge theory superpotentials is to use the matrix model to compute the effective superpotential in the “deepest infrared” in the Wilsonian sense, and then to integrate in all these fields. This seems to limit the use of matrix models in describing theories which one cannot obtain by integrating in fields.

This procedure of reconstructing the gauge theory superpotential out of the matrix model results may seem somewhat cumbersome. Recently, for theories with tree-level superpotential expressed only in terms of quark bilinears (mesons), a method was proposed [13] to link the matrix model and the gauge theory directly, bypassing the detour through the “deepest infrared”. The method consists of introducing in the matrix path integral a δ function which relates the matrix model quark bilinear with the gauge theory meson. While this method seems rather ad hoc, as it identifies two quantities which have nothing to do with each other (they are related by a dictionary, but they cannot, in principle, be identified), it implies that the full effective superpotential is given by the sum between the tree level superpotential and an Affleck-Dine-Seiberg nonperturbatively-generated term. The fact that this is the correct answer can be argued using standard gauge theory arguments.

It is unclear at this time how to extend this program to directly include superpotentials with terms depending of several types of fields (e.g. Yukawa coupling), with terms which cannot be written in terms of quark bilinears (e.g. baryon operators in SQCD with $N_f \geq N_c$).

This procedure is moreover not unique. In the appendix we present an alternative procedure relating the matrix model and the gauge theory which also gives the correct gauge theory result and moreover treats different representations on the same footing. Both the procedure of [13] as well as the one described in the appendix are best thought of as *shortcuts* which relate the two sides before integrating out the fields on either side. It would be quite interesting if one can justify them to be more than this.

Let us proceed by reviewing the method proposal of [13]. The idea is to identify the matrix model gauge invariant quark bilinear with the field theory meson, by introducing a constraint for the quarks in the matrix integral. More formally,

$$e^{-\mathcal{F}_{\xi=1}} = \int DQ_i D\tilde{Q}_i \delta(M_{ij} - Q_i \tilde{Q}_j) e^{-W_{tree}(Q_i \tilde{Q}_i)} \quad , \quad (17)$$

where the δ function enforces a direct relation between the high energy description (in terms of the independent fields Q_i^α and $\tilde{Q}_\alpha^i$) and the one in terms of composite fields M_{ij} .

Because of this δ function, any tree level superpotential which can be expressed as a function of the mesons automatically gets out of the integral. The remainder of the integral gives the volume of the constrained quark integration domain. As in the original DV prescription, the glueball superfield is identified with the dimension of the matrix model matrices (at least in the case of $U(M)$ matrix model gauge group) after it was taken to be large. Adding the usual Veneziano-Yankielowicz terms (to account for the gauge field dynamics [15]) and integrating out the glueball superfield one obtains the Affleck-Dine-Seiberg superpotential. The net result of (17) is to always give the meson effective potential

$$W_{eff} = W_{ADS} + W_{tree} \quad . \quad (18)$$

For simple tree level superpotentials this is indeed what one expects to find in gauge theory [31]. However, it is not a priori clear that for a generic tree level superpotential the ADS superpotential is the only possible nonperturbative contribution. If this were not the case, the procedure (17) would fail.

We can outline the sketch of a gauge theoretic argument that indeed there is no nonperturbative contribution besides the ADS piece. Let us assume a W_{tree} of the form

$$W_{tree} = \sum_i g_{i,k} Tr[X^i]^k \quad (19)$$

Besides the ADS superpotential, one can expect functions of the dimensionless and chargeless ratios of coupling constants (e.g. $\frac{g_{1g4}}{g_{2g3}}, \frac{g_{1g5}}{g_{2g4}}, \frac{g_{2g17}}{g_{12g7}}$, etc.) to appear multiplying W_{ADS} . However, these functions need to be expandable in positive integer powers of the coupling constants around the origin. Moreover they need to be holomorphic in all couplings. Hence, these functions

would be holomorphic functions which go to zero at ∞ , and would thus be zero everywhere. Thus, the ADS superpotential will be the only nonperturbative contribution to the superpotential.

As pointed out in the beginning, the procedure (17) is an effective way to bypass integrating out on both sides when the potential can be expressed in terms of mesons only. However, extending it to fields in other representations appears challenging.

For example, in theories containing massless adjoints the number of mesons one can construct becomes very large [32, 33]. Multiple mesons can appear also in the case one considers theories with massless quarks in the symmetric or the antisymmetric representations of the gauge group. In order to capture the dynamics of these mesons, one would need to add them by hand in (17). This would further constrain the quark integration volume. As before, any superpotential depending on the mesons would get out of the integral, and the constrained volume would give the nonperturbative superpotential as a function of all these mesons.

Nevertheless, if one tries to apply this procedure to baryons there is no way to get sensible results. Indeed, a gauge theory baryon is a combination of N_c quarks. However, the matrix model prescription instructs us to take the matrix model gauge group to have large rank, unrelated to the gauge theory N_c . Thus, depending upon how one chooses to treat it, the naive baryon operator either vanishes or is not gauge invariant. A matrix theory baryon should be composed of M quarks, where M is the dimension of the matrix. It is quite clear that the matrix model baryon and the gauge theory one have nothing in common with each other.

One can actually see that baryons are a problem not only for various computational “shortcuts”, but also for formulating the correspondence in the picture in which one first goes to the deep infrared of both theories. One should add a term of the form $b\epsilon^{i_1,\dots,i_{N_c}}\epsilon_{j_1,\dots,j_{N_f}}Q_{i_1}^{j_1}\dots Q_{i_{N_c}}^{j_{N_f}}$ in the gauge theory and integrate it out. Nevertheless, it is quite clear that such a term does not make any sense in the large M $U(M)$ matrix model where the quarks transform in the fundamental representation of $U(M)$ but their number remains fixed to N_f . The next section is devoted to solving this problem.

4 Baryons

In this section we try to understand how baryons enter in the gauge theory - matrix model correspondence. As we explained above, introducing baryon sources *both* in the large M matrix model potential and in the gauge theory superpotential cannot be done. Thus, one has to take a step back and consider the motivation behind the DV proposal.

The proposal was first formulated via a string theory route. Unfortunately, understanding the role of fundamentals from this standpoint is rather challenging. In the last section of this paper we will return to the question of how one can understand theories with fields in the adjoint and fundamental representations via geometrical transition.

The purely gauge theoretical motivation of this duality was formulated in [4], and in a nutshell can be summarized as: “planar diagrams in theories with adjoints compute holomorphic quantities.”

If one adopts this outlook, the matrix model is merely a tool to systematically compute planar diagrams. Phrased in this way, the “dictionary” between the two sides does nothing but select the planar diagrams in the gauge theory. Of course, it is still amazing that planar diagrams compute quantities which one normally obtains nonperturbatively, but the “miracle” now pertains only to the gauge theory side.

The extension to theories with fundamentals and no baryons was conjectured in [12] to be: “planar diagrams and diagrams with one quark boundary compute holomorphic quantities”. In section 2 we gave a gauge theoretical proof, along the lines of [15], that indeed only diagrams with one boundary contribute to the low energy effective superpotential. As a side remark, we should note that there is still no string theory motivation for this extension.

In the same spirit as [4], the matrix model with quarks and the correct dictionary between the two sides [9, 12] should be thought of only as a tool to systematize the gauge theory computation of these diagrams with boundaries.

If one introduces now diagrams with baryons, it is clear that they are quite different from diagrams with quark loops. Indeed, the quark lines are not closed, but rather go from the baryons to the antibaryons. Nevertheless, we have shown in section 2 that a pair of baryons can be thought of as a multi-trace operator. We have also introduced a generalized notion of boundary for this kind of operators, and we have proven that only planar Feynman diagrams with one generalized boundary can generate the effective

superpotential.

The aim is therefore to analyze in a systematic way these planar diagrams. Unfortunately, a large M matrix model appears not to be any more the right “tool” for selecting these diagrams. Baryon terms of the form $b\epsilon^{i_1,\dots,i_{N_c}}\epsilon_{j_1,\dots,j_{N_f}}Q_{i_1}^{j_1}\dots Q_{i_{N_c}}^{j_{N_f}}$ cannot be introduced in the Matrix Lagrangian, because the matrix quarks are $1 \times M$ matrices, and the combinations above are not matrix scalars.

Thus, the only types of gauge theory superpotentials one can insert in the large M Matrix Lagrangian are the ones which can be expressed in terms of color singlet quark bilinears. As we have seen in the previous section, in this case the shortcut of [13] can be applied, and one simply obtains the sum of the ADS superpotential and the tree level one.

Since the large M matrix model cannot be used as a tool to select planar Feynman diagrams when baryons are present, the only way to count these diagrams is to select them by hand in a matrix model whose gauge group $U(M)$ has the same rank as the gauge theory gauge group $U(N_c)$. We stress that this complication is necessary only when baryons (or more generally operators which exist only for fixed values of the N_c) are present.

By identifying again g^2M with S , one obtains an effective superpotential $W_{eff}(S, m_i, b, \tilde{b} \dots)$. Counting these diagrams is quite nontrivial, especially for baryons with many quarks, or for theories containing other terms besides quark masses and baryons.

If one integrates S out of the matrix result $W_{eff}(S, m_i, b, \tilde{b} \dots)$, and integrates in the quarks one should obtain the full gauge theory result.

It is also possible to see that, due to charge conservation, a vacuum diagram contains an equal number of baryons and anti-baryons. The product of a baryon and an antibaryon can be reexpressed as a multiple trace operator, by using the fact that a product of ϵ tensors can be expressed as a sum of products of Kronecker δ functions. Therefore, any nonzero vacuum Feynman diagram containing baryons can be written as a diagram containing multi-trace operators, and as such the large M limit gives sensible results.

Alternatively, one can integrate out a quark linking the two baryon, and obtain an effective di-baryon operator which again makes sense in the large M limit.

It is not hard to see that this discussion combined with the fact that multi-trace tree level potentials in the gauge theory only receive W_{ADS} as a non-perturbative correction extend the correspondence to theories with baryons.

In the next section we will compute the effective superpotential for an

$U(2)$ gauge theory with 2 flavors, and show that one recovers the expected gauge theory result. We then use the shortcut (17) and recover the quantum moduli space of $SU(N)$ SQCD with N flavors.

5 Effective superpotentials for mesons and baryons from Matrix Models

After discussing the mesons and the baryons, we now turn to calculating the gauge theory superpotential from matrix model. We start with a brief review of the field theory results for theories where $\mathcal{N} = 2$ SUSY is broken to $\mathcal{N} = 1$.

5.1 Field Theory results

Consider⁴ $\mathcal{N} = 1$ $SU(N_c)$ theories with N_f quarks in the fundamental representation, obtained from $\mathcal{N} = 2$ $SU(N_c)$ by turning on a mass term for the adjoint chiral multiplet:

$$W = \sqrt{2}\text{Tr}(\tilde{Q}\phi Q) + \mu\text{Tr}\phi^2 \quad (20)$$

To discuss the moduli space of this theory, we distinguish between the case when the mass of the adjoint field is small (the adjoint field remains as a field in the $\mathcal{N} = 1$ theory) and the case when the mass of the adjoint field is large and the adjoint is integrated out and we obtain $\mathcal{N} = 1$ supersymmetric QCD with the dynamical scale:

$$\Lambda_{\mathcal{N}=1}^{3N_c-N_f} = \mu^{N_c} \Lambda_{\mathcal{N}=2}^{2N_c-N_f} \quad (21)$$

After integrating out the adjoint field, one obtains:

$$W = \frac{1}{2\mu} \left[\text{Tr}[(Q\tilde{Q})(Q\tilde{Q})] - \frac{1}{N_c} \text{Tr}(Q\tilde{Q})\text{Tr}(Q\tilde{Q}) \right] \quad (22)$$

For $\mu \rightarrow \infty$, this superpotential approaches zero and we then obtain $\mathcal{N} = 1$ SQCD.

⁴Unlike the rest of the paper, in this section we consider the gauge group to be $SU(N_c)$ rather than $U(N_c)$. We do this to conform with the papers referred to in this section. All relevant results also hold for $U(N_c)$.

In [34, 35], a detailed discussion was presented, relating the vacua of $\mathcal{N} = 2$ SUSY and $\mathcal{N} = 1$ SUSY theories. Without going into details, we recall now the main results regarding the $\mathcal{N} = 1$ vacua:

- for $N_f \leq N_c - 1$, the strong gauge dynamics corrects the tree level superpotential by the addition of the ADS superpotential (22). The moduli space is the extrema of this superpotential and, at finite values for the mass of the adjoint field, it has been shown to have 2 types of solutions [35], one when all the diagonal entries of $M = Q\tilde{Q}$ are equal to

$$M_{ii} = \left(\frac{N_c}{N_c - N_f}\right)^{\frac{N_c - N_f}{2N_c - N_f}} \mu \Lambda \quad (23)$$

and the other one when there are two different diagonal entries (there are only two possible values for the diagonal entries). The presence of this two choices remains valid if we add masses for the quarks and integrate them out. The first case has been discussed in matrix model in [9, 12]. The second one can be treated similarly.

In a pure $N = 1$ SQCD, with vanishing tree level superpotential, all these vacua run away to infinity.

- for $N_f = N_c$, the situations changes because the classical moduli space of vacua is not only parametrized by mesons, but also by baryons. In the case $N_f = N_c$, the classical moduli space of vacua is modified quantum mechanically by

$$\det M - B\tilde{B} = \Lambda^{2N_c} \quad (24)$$

In this case there is no nonperturbatively-generated correction to the effective superpotential; the constraint (24) is implemented with a Lagrange multiplier X . The exact effective superpotential is then

$$W_{eff} = W = X(\det M - B\tilde{B} - \Lambda^{2N_c}) + \frac{1}{2\mu} \left[\text{Tr}(M^2) - \frac{1}{N_c} \text{Tr}(M) \text{Tr}(M) \right] \quad (25)$$

As before, M can again have at most two different eigenvalues; for $X = 0$ we return to the situation discussed before, for $X \neq 0$ one has new branches which did not exist for $N_f \leq N_c - 1$:

$$M_{ii} = \Lambda^{2N_c} \quad (26)$$

for $B\tilde{B} = 0$ or

$$M_{ii}^{N_c} - B\tilde{B} = \Lambda^{2N_c} \quad (27)$$

for $B\tilde{B} \neq 0$

- for $N_f \geq N_c + 1$, the classical moduli space is not modified, but the number of baryons increases. In this range, there is a dual magnetic description based on the gauge group $SU(N_f - N_c)$ with N_f fundamental flavors and gauge invariant fields M , the Seiberg-dual theory [36]. Discussions on the realization of Seiberg's duality for mass deformed theories in the context of gauge theory-matrix models appeared in [17], but it is unclear how to obtain a duality for massless theories.

5.2 Quantum Moduli Spaces from Matrix Models

As explained earlier, the fields appearing in the matrix model are not directly related to the dynamical gauge theory fields. Thus, one cannot directly identify the matrix model effective action in which only some fields were integrated out with the gauge theory superpotential. Instead, to find the gauge theory effective superpotential using the matrix model, we take the following steps:

- 1) First we deform the gauge theory by including mass terms for all fields as well as sources for the operators describing the directions in the moduli space we are interested in. In the case of mesons these two coincide, as the mass of a quark acts as a source for a meson as well.

- 2) We compute the planar free energy of the matrix model associated to this deformed theory.

- 3) We equate the resulting function with the gauge theory effective superpotential at energy scales below the lightest mass present in the deformed theory. The 't Hooft coupling of the matrix model is identified with the gauge theory glueball superfield. Furthermore, the Veneziano-Yankielowicz superpotential is added to take into account the gauge dynamics.

- 4) We integrate in gauge theory fields by appropriately performing Legendre transforms

- 5) We restore the masses of the original fields, by appropriately taking the deformation parameters to zero.

To illustrate this program let us discuss in detail an $SU(2)$ gauge theory with 2 flavors, Q_i and $\tilde{Q}_i$ with $i = 1, 2$. This means that the adjoint field (appearing in the discussion in the previous subsection) has been integrated out and we work within $\mathcal{N} = 1$ SQCD. We return at the end of this section to the case when the adjoint field has finite mass. We start with a massless theory whose tree level superpotential is given by the baryon and antibaryon

sources:

$$W_{tree} = \int d^2\theta \left[bQ_1^\alpha Q_2^\beta \epsilon_{\alpha\beta} + \tilde{b}\tilde{Q}_1^\alpha \tilde{Q}_2^\beta \epsilon_{\alpha\beta} \right] + hc \quad (28)$$

The first step then instructs us to define this theory as the massless limit of the theory with the superpotential above deformed by mass terms:

$$W_{deff} = \int d^2\theta \left[mQ_1^\alpha \tilde{Q}_\alpha^1 + mQ_2^\alpha \tilde{Q}_\alpha^2 \right] + hc \quad (29)$$

As discussed earlier, it is clear that W_{tree} cannot be used as part of the potential of a large N matrix model (or any other theory for that matter), because it manifestly vanishes. Thus, the way to proceed is to consider an $SU(2)$ matrix model and compute only the planar free energy. Thus, we have the following matrix model potential:

$$W_M = (Q_1^\alpha, Q_2^\alpha, \tilde{Q}_\alpha^1, \tilde{Q}_\alpha^2) \begin{pmatrix} 0 & b\epsilon_{\alpha\beta} & m & 0 \\ b\epsilon_{\alpha\beta} & 0 & 0 & m \\ m & 0 & 0 & \tilde{b}\epsilon^{\alpha\beta} \\ 0 & m & \tilde{b}\epsilon^{\alpha\beta} & 0 \end{pmatrix} \begin{pmatrix} Q_1^\beta \\ Q_2^\beta \\ \tilde{Q}_\beta^1 \\ \tilde{Q}_\beta^2 \end{pmatrix} \equiv \hat{Q}^T \hat{M} \hat{Q} \quad (30)$$

Since this potential is quadratic in the dynamical fields Q and $\tilde{Q}$, the free energy is trivial to compute. The result is:

$$\mathcal{F} = -\ln Z = \ln \left[\frac{1}{\Lambda^{2 \times 2}} \det \hat{M} \right]^{1/2} = 2 \times 2 \times \frac{1}{2} \ln \left(\frac{m^2 - b\tilde{b}}{\Lambda^2} \right) \quad (31)$$

In the expression above the first factor of 2 is due to having two flavors while the second factor of 2 is due to having two colors. This latter numerical coefficient combines with the coupling constant into the glueball superfield. Therefore the gauge theory superpotential is:

$$W_G = N_c S \left[1 - \ln \frac{S}{\Lambda^3} \right] + \frac{1}{2} N_f S \ln \left(\frac{-b\tilde{b} + m^2}{\Lambda^2} \right) \quad (32)$$

where in the case at hand $N_f = N_c = 2$ but we wrote them explicitly to stress the functional dependence, and we inserted Λ for dimensional reasons.

Now we perform the Legendre transform required for integrating in the quarks in the form of the gauge theory mesons $X_i = Q_i \tilde{Q}^i$ and baryons $B = Q_1^\alpha Q_2^\beta \epsilon_{\alpha\beta}$ and $\tilde{B} = \tilde{Q}_1^\alpha \tilde{Q}_2^\beta \epsilon_{\alpha\beta}$. This implies that we deform W_G by

$$W_L = -mN_f X - bB - \tilde{b}\tilde{B} \quad (33)$$

and integrate out the sources m , b and $\tilde{b}$. Here we also considered all mesons to be identical. Eliminating the sources by their equations of motion and using the equality between N_c and N_f we find:

$$W_{eff}(X, B, \tilde{B}, S) = 2S \ln \frac{\Lambda^4}{-B\tilde{B} + X^2} . \quad (34)$$

Upon eliminating S by its equation of motion we find the correct gauge theory superpotential

$$W_{eff} = 0 \quad (35)$$

together with the quantum corrected moduli space:

$$\det X - B\tilde{B} = \Lambda^4 \quad (36)$$

where we wrote X^2 in the $SU(2) \times SU(2)$ invariant form. This reproduces the gauge theory result.

5.3 The Effective Potential from Integrating Out the Massive Quarks

Let us now consider the general case of an $SU(N_c)$ theory with $N_f = N_c$ flavors and discuss the relation between the matrix model and gauge theory. Since we are interested in incorporating baryonic operators in the matrix model, we deform the gauge theory action by source terms for them. Then, the idea we will pursue is the following. Using matrix model reasoning we reduce the matrix model with $N_f = N_c$ flavors to one with $N_f = N_c - 1$ flavors. Then, for this new matrix model we construct the corresponding gauge theory superpotential. Then we integrate in, as gauge theory quark, the quark which was integrated out as a matrix model one as well as the baryons. To make a long story short, we will recover the known gauge theory result, i.e. a vanishing effective superpotential and a quantum-deformed moduli space.

To begin with, the deformed tree level superpotential is:

$$W_{def} = mQ_1^\alpha \tilde{Q}_{1\alpha} + bB + \tilde{b}\tilde{B} \quad (37)$$

where the baryon operators B and $\tilde{B}$ are defined in equation (12). Since B and $\tilde{B}$ are antisymmetric products of the N_f quarks, we can split them into products of the massive quark and the remaining massless quarks:

$$B = Q_{1\alpha} \hat{B}^\alpha \quad (38)$$

where $\hat{B}^\alpha$ is product of $Q_2, \dots, Q_{N_c}$. The same applies for the antibaryon:

$$\tilde{B} = \tilde{Q}_{1\alpha} \hat{\tilde{B}}^\alpha \quad (39)$$

With these slight adjustments, the superpotential reads

$$W_{def} = mQ_1\tilde{Q}_1 + bQ_{1\alpha}\hat{B}^\alpha + \tilde{b}\tilde{Q}_{1\alpha}\hat{\tilde{B}}^\alpha \quad (40)$$

Since the massive quarks appear only linearly and quadratically, it is easy to compute the matrix model effective action by integrating out Q_1 and $\tilde{Q}_1$ as matrix model fields. The computation is entirely straightforward and yields

$$W_{eff}^{massive} = S \ln \frac{m}{\Lambda} - \frac{b\tilde{b}}{4m} (\hat{B}^\alpha \hat{\tilde{B}}_\alpha) \quad (41)$$

where we identified the size of matrices in the matrix model with the gauge theory glueball superfield.

Because $\hat{B}^\alpha$ and $\hat{\tilde{B}}^\alpha$ are products of $N_c - 1$ flavors,

$$\hat{B}^\alpha \hat{\tilde{B}}_\alpha = \det(M_{N_f=N_c-1}) \quad (42)$$

To obtain the gauge theory superpotential one should also integrate over the massless quarks. Instead of computing the free energy with this effective potential and then integrating in the gauge theory quarks, we can just apply the prescription [13] which we reviewed in section 3. This straightforwardly gives:

$$\begin{aligned} W_{eff}(S, M_{N_c-1}, m, b, \tilde{b}) &= S \left(-\ln \frac{S}{\Lambda^3} + 1 \right) - S \ln \left[\frac{\det(M_{N_f=N_c-1})}{\Lambda_{N_f=N_c}^{2N_c-1}} \right] \\ &+ S \ln \frac{m}{\Lambda} - \frac{b\tilde{b}}{4m} (\hat{B}^\alpha \hat{\tilde{B}}_\alpha) \quad (43) \end{aligned}$$

The term proportional of the logarithm of the mass of the quark which was integrated out combines with the initial dynamical scale Λ to give the scale of the theory with $N_f = N_c - 1$ quarks, just as in a purely field-theoretic analysis. Then, to construct the gauge theory superpotential in terms of meson fields only, we integrate out the glueball superfield and find the usual ADS superpotential:

$$W_{eff}(M_{N_c-1}) = \frac{\Lambda_{N_f=N_c-1}^{2N_c+1}}{\det(M_{N_f=N_c-1})} - \frac{b\tilde{b}}{4m} (\hat{B}^\alpha \hat{\tilde{B}}_\alpha) \quad (44)$$

The last step is to integrate in the gauge theory quark corresponding to the matrix model quark which was integrated out, and restore the last meson as well as the two baryons. The steps are identical to those in [30] and we will not repeat them here. The conclusion of this short computation is that we recover a vanishing superpotential as well as the usual modification of the classical moduli space of the theory.

6 Geometric Transitions, Matrix Models and Gauge Theories for Fundamental Quarks

The geometric transition was first formulated in type IIA string theory from one side with D6 branes wrapped on the $\mathbf{S}^3$ cycle of a deformed conifold to another side with fluxes on the $\mathbf{P}^1$ cycle of a resolved conifold [37] (see [38, 39, 40, 46] for subsequent work on type IIA).

The mirror picture is a type IIB transition from D5 branes wrapped on the $\mathbf{P}^1$ of a resolved conifold to fluxes on a $\mathbf{S}^3$ cycle of the deformed conifold. The type IIB picture was easier to generalize, by using $\mathcal{N} = 1$ deformations of resolved $\mathcal{N} = 2$ ADE singularities [41, 42, 43, 44, 45].

The geometrical picture was the backbone of the recent exciting developments in matrix model/gauge theories. In [1], the model of [41] has been used to relate matrix model variables with geometrical variables and with gauge theory variables. In the comparison between geometry, matrix models and gauge theories, what needs to be compared is:

- in field theory we consider the quantum moduli space. One of the tools to study this are the Seiberg-Witten curves. Going beyond the field theory, gravitational corrections to field theory have been discussed in the matrix model description [47, 48].
- in geometry, we consider the 1-complex dimensional curve obtained by reducing the deformed geometry on an S^2 fiber.
- in the matrix model, we consider the equation for the resolvent of the matrix.

If we are to compare just field theory and matrix model, in [3] has been suggested that the planar diagrams in matrix models give the non-perturbative corrections in the field theory (we have discussed extensively in our paper about this). In the line of [1], it is interesting to get a connection between field theory and matrix model, through geometrical quantities. It

would be interesting to also describe matter (besides bifundamental fields) in the geometrical setting. It is not yet known how to go through the transition with fundamental matter, and in the present section we will outline a possibility, leaving the details for a future work.

To begin with, we have to know how to include fundamental matter in the geometrical set-up. The fundamental matter would correspond to either additional D5 branes wrapped on additional non-compact cycles of the conifold or to additional D7 branes which are partially wrapped on the $\mathbf{P}^1$ cycles. In order to see this, one can use the MQCD picture for the geometrical transition [49, 50, 51, 52]. In the absence of the fundamental quarks, the configuration with D5 branes wrapped on $\mathbf{P}^1$ cycles goes into a brane configuration with D4 branes between two orthogonal NS branes [53] (there are two orthogonal lines of singularity in the geometry, which become two orthogonal NS branes after the T-duality). In the brane configuration, the fundamental matter can be introduced as either semi-infinite D4 branes ending on one of the NS branes or as D6 branes in between the NS branes. The D4 branes are better suited to describe massive flavors and the D6 branes are better suited to describe massless flavors. In the geometric picture the semi-infinite D4 branes become D5 branes on a non-compact $\mathbf{P}^1$ which is distanced from the exceptional $\mathbf{P}^1$ by a distance equal to the mass of the flavor [50]. We do not yet have a clear picture on the geometry side for the D6 brane, but we can use the geometry/brane configuration duality in order to discuss the transition.

For massive matter, in the infrared we go to the effective theory where all the massive matter has been integrated out and the sizes of the different $\mathbf{S}^3$ cycles on the geometric side are related to the different gluino condensates in field theory, which are in turn proportional to the masses of the quarks. If one of the masses of the quarks becomes zero, the size of the $\mathbf{S}^3$ cycle becomes zero and the geometric transition fails. In this case, we should use the brane configuration with D6 branes instead of the one with semi-infinite D4 branes. By lifting it to MQCD and going through the transition, the moduli space of the field theory should be recovered from the geometry. As the transition consists on closing the interval between the two NS branes (going from a curved M5 brane to a plane M5 brane [49, 50]), we expect to get a geometry described by an equation $xy = a$ in the (x, y) plane. But section 5.2 of [35] tells us that the only case when $xy \neq 0$ is when the mass of the $\mathcal{N} = 2$ adjoint field is not infinite. This is consistent with the fact that there is no supersymmetric vacua for the corresponding $\mathcal{N} = 1$ theory.

In the case of infinite mass for the adjoint field, after integrating it out in the matrix theory, we remain with decoupled integrals over the massless fundamental quarks (with no tree level superpotential), and this gives the ADS superpotential which removes any supersymmetric vacua.

In order to obtain a vacuum, we have to keep the mass of the $\mathcal{N} = 2$ adjoint field as a finite quantity. In this case the matrix integral becomes harder to compute, as there are extra couplings between fields. This is currently under investigation.

We can ask whether there exists a geometrical picture corresponding to finite mass for the $\mathcal{N} = 2$ adjoint field. Remember that the starting point of the geometrical discussion was the case when the mass of the $\mathcal{N} = 2$ adjoint field is ∞ . In the brane configuration this comes from the fact that the NS branes are orthogonal and this comes from the orthogonality of two lines of singularity in the geometry. As we know from brane configurations, the angle between the NS branes is related to the mass of the adjoint, therefore a finite mass for the adjoint field can be obtained only when the NS branes are not orthogonal which would imply a geometry with non-orthogonal lines of singularity.

7 Conclusions

In this paper we discussed several aspects of the Dijkgraaf-Vafa relation between matrix models and gauge theories. Our investigations focused on the fields transforming in the fundamental representation of the gauge group. Using field theory arguments (based on symmetries and holomorphy) we proved that the effective superpotential for the glueball superfield is generated on the gauge theory side by planar (in a slightly generalized sense) Feynman diagrams with exactly one boundary. Our result holds for a tree level superpotential given by a generic polynomial in multi-trace operators and baryonic couplings. It is then likely that, using techniques similar to those in [4], the gauge theory computation be explicitly reduced to a matrix model computation. A slight subtlety in the diagrammatic analysis is due to the existence of graphs which are not obviously planar, but can be mapped into manifestly planar ones using the symmetries of the baryonic vertices.

We also analyzed the inclusion of baryonic sources in the matrix model computation. This solution turned out to be rather subtle, as one cannot include the baryon operators in a large N matrix model computation. In-

stead one must keep the rank of the matrix model gauge group equal to the rank of the gauge theory one. Then, the restriction to planar diagrams is not automatic any more and must be done by hand. We explicitly performed the computations for $U(2)$ SQCD with two flavors and then used a “computational shortcut” to analyze the general case of $U(N)$ SQCD with N flavors. We recovered the known gauge results for the quantum moduli space.

As the geometry was a crucial tool in deriving the relation between matrix models and gauge theories in [1], it remains an interesting open problem finding the deformations of the geometry which correspond to the addition of fundamental matter to the gauge theory. We have outlined a possible way to describe the required geometrical transition. We plan to return to this in a future work.

Appendix

In this appendix, using heuristic arguments, we propose a computational shortcut in the matrix model which yields the gauge theory effective superpotential.

The proposal implies that the effective superpotential is written as a sum of terms, each arising from one particular field in the theory. We will be concerned only with the contribution of matter fields. As in the usual DV approach, the gauge field dynamics is taken into account by the addition of the Veneziano-Yankielowicz superpotential W_{VY} to the matrix model free energy.

Our proposal is that the free energy of the matrix model associated to some gauge theory is given by the logarithm of its non-normalized partition function. Then, the gauge theory effective superpotential is given by the (extended) DV relation:

$$W_{effective} = W_{VY} + \frac{\partial \mathcal{F}_{\chi=2}}{\partial S} + \mathcal{F}_{\chi=1} \quad . \quad (45)$$

An immediate consequence of this proposal is that a field not appearing in the superpotential contributes a factor of its “integration volume”:

$$\int D\Psi_i = \text{Vol}(\Psi_i) \quad . \quad (46)$$

As usual, we identify the 't Hooft coupling in the matrix model, $g^2 M$, (where M is dimension of the matrices) with the glueball superfield S .

In the case of massless adjoints, their volume of integration is the volume of $SU(M)$ multiplied by that of a noncompact piece. When properly regularized the volume of the noncompact piece becomes an arbitrary scale. This contribution comes in the numerator, and can cancel the “volume of gauge group” which gives the VY term. If the gauge theory was $N = 2$ supersymmetric, there should not be any superpotential⁵ for the glueball superfield S . This cancellation can be achieved if one identifies the cutoff in the volume of the adjoint field with the dynamically generated scale, Λ .

Let us consider now the case of massless fields in the fundamental representation. Their integration volume is also infinite, and one can also handle it by imposing a cutoff. However, it is not *a priori* clear what this cutoff should be identified with in the gauge theory. It is nevertheless clear that one should use an object invariant under all symmetries of the (gauge) theory which also has the right dimension. A natural choice seems to be the determinant of the meson field:

$$\det X = |Q\tilde{Q}|_{max} \quad . \quad (47)$$

To compute the integration volume for one quark we notice that the measure $dQ d\tilde{Q}$ spans two fundamental representation; their product can be thought of as giving an adjoint representation and a singlet. The volume of the adjoint is finite, but the volume of the singlet is infinite. If we regularize this volume by imposing a cutoff at a “radius” X , the total volume will just be volume of the adjoint multiplied by X^{M^2} ⁶.

For each of the N_f quarks not appearing in the superpotential, one has to consider the corresponding integration volume. The cutoff for each volume will be interpreted in the gauge theory as the meson corresponding to that quark. Due to the various global symmetries of the theory these cutoffs should combine to form an invariant, e.g. $\det X$. It is not hard to see that, the net result of these considerations is a $W_{eff}(S, X_{ij})$ which yields the Affleck-Dine-Seiberg superpotential upon integrating out the glueball superfield.

Before integrating out the glueball superfield, the superpotential can be

⁵The full analysis of the $N = 2$ dynamics is far beyond the scope of this proposal, see [27] for steps in that directions.

⁶This can be easily understood by analogy with the volume of a ball. The ball can be thought as a product of a sphere and R^+ . The total volume is the product of the volume of the sphere and the radius to a power equal to the dimension of the space.

justified to be given by:

$$W = N_c S \left[\log \left(\frac{\Lambda^3}{S} \right) + 1 \right] - N_f S \left[\log \left(\frac{\Lambda^3}{S} \right) + 1 \right] - S \log \left(\frac{\det X_{ij}}{\Lambda^{2N_f}} \right) \quad (48)$$

Indeed, the first term is the contribution of the gauge field; the second term comes from the compact part of the integration volume of the quarks; finally, the last term is given by the non-compact directions in the quark integration volumes, combined in a way compatible with the global symmetries of the theory.

The mesons X_i appear now as cutoffs (regulators) in the integration volumes, and each gives a contribution to $\mathcal{F}$ of the form $S^2 \log X_i$, as explained above. This justifies the X dependence of (48), as the superpotential is related to the first derivative of $\mathcal{F}$ with respect to S ,

When $N_f = N_c$, the first two terms cancel, and the only part left is $S \log \frac{\det X}{\Lambda^{2N_c}}$. Integrating out S gives a vanishing superpotential as well as the constraint

$$\Lambda^{2N_f} = \det X \quad (49)$$

which is the correct gauge theory result in the absence of baryons.

Acknowledgments

We thank Eric d'Hoker, Per Kraus and Hitoshi Murayama for useful discussions. The work of I.B. was supported by the NSF under Grant No. PHY00-99590. The work of R.R. was supported in part by DOE under Grant No. 91ER40618 and in part by the NSF under Grant No. PHY00-98395. R.T. was supported by the DOE grant DE-AC03-76SF00098, the NSF grant PHY-0098840 and by the Berkeley Center for Theoretical Physics.

References

- [1] R. Dijkgraaf and C. Vafa, “Matrix models, topological strings, and supersymmetric gauge theories,” Nucl. Phys. B **644**, 3 (2002), hep-th/0206255.
- [2] R. Dijkgraaf and C. Vafa, “On geometry and matrix models,” Nucl. Phys. B **644**, 21 (2002), hep-th/0207106.

- [3] R. Dijkgraaf and C. Vafa, “A perturbative window into non-perturbative physics,” hep-th/0208048.
- [4] R. Dijkgraaf, M. T. Grisaru, C. S. Lam, C. Vafa and D. Zanon, “Perturbative computation of glueball superpotentials,” hep-th/0211017.
- [5] H. Fuji and Y. Ookouchi, “Comments on effective superpotentials via matrix models,” hep-th/0210148.
- [6] D. Berenstein, “Quantum moduli spaces from matrix models,” hep-th/0210183.
- [7] R. Dijkgraaf, S. Gukov, V. A. Kazakov and C. Vafa, “Perturbative analysis of gauged matrix models,” hep-th/0210238.
- [8] A. Gorsky, “Konishi anomaly and $N = 1$ effective superpotentials from matrix models,” hep-th/0210281.
- [9] R. Argurio, V. L. Campos, G. Ferretti and R. Heise, “Exact superpotentials for theories with flavors via a matrix integral,” hep-th/0210291.
- [10] J. McGreevy, “Adding flavor to Dijkgraaf-Vafa,” hep-th/0211009.
- [11] H. Suzuki, “Perturbative derivation of exact superpotential for meson fields from matrix theories with one flavour,” hep-th/0211052.
- [12] I. Bena and R. Roiban, “Exact superpotentials in $N = 1$ theories with flavor and their matrix model formulation,” hep-th/0211075.
- [13] Y. Demasure and R. A. Janik, “Effective matter superpotentials from Wishart random matrices,” hep-th/0211082.
- [14] R. Gopakumar, “ $N = 1$ theories and a geometric master field,” hep-th/0211100.
- [15] F. Cachazo, M. R. Douglas, N. Seiberg and E. Witten, “Chiral Rings and Anomalies in Supersymmetric Gauge Theory,” arXiv:hep-th/0211170.
- [16] R. Dijkgraaf, A. Neitzke and C. Vafa, “Large N strong coupling dynamics in non-supersymmetric orbifold field theories,” arXiv:hep-th/0211194.

- [17] B. Feng, “Seiberg Duality in Matrix Model”, hep-th/0211202, B. Feng and Y. H. He, “Seiberg Duality in Matrix Models II,” arXiv:hep-th/0211234.
- [18] N. Dorey, T. J. Hollowood and S. P. Kumar, “S-duality of the Leigh-Strassler deformation via matrix models,” arXiv:hep-th/0210239. N. Dorey, T. J. Hollowood, S. P. Kumar and A. Sinkovics, “Massive vacua of $N = 1^*$ theory and S-duality from matrix models,” arXiv:hep-th/0209099. N. Dorey, T. J. Hollowood, S. Prem Kumar and A. Sinkovics, “Exact superpotentials from matrix models,” arXiv:hep-th/0209089.
- [19] M. Aganagic, A. Klemm, M. Marino and C. Vafa, “Matrix model as a mirror of Chern-Simons theory,” arXiv:hep-th/0211098.
- [20] F. Ferrari, “Quantum parameter space and double scaling limits in $N = 1$ super Yang-Mills theory,” arXiv:hep-th/0211069. F. Ferrari, “On exact superpotentials in confining vacua,” arXiv:hep-th/0210135.
- [21] H. Fuji and Y. Ookouchi, “Comments on effective superpotentials via matrix models,” arXiv:hep-th/0210148.
- [22] D. Berenstein, “Quantum moduli spaces from matrix models,” arXiv:hep-th/0210183.
- [23] R. Dijkgraaf, S. Gukov, V. A. Kazakov and C. Vafa, “Perturbative analysis of gauged matrix models,” arXiv:hep-th/0210238.
- [24] S. G. Naculich, H. J. Schnitzer and N. Wyllard, “The $N = 2$ $U(N)$ gauge theory prepotential and periods from a perturbative matrix model calculation,” arXiv:hep-th/0211123.
- [25] R. Argurio, V. L. Campos, G. Ferretti and R. Heise, “Baryonic Corrections to Superpotentials from Perturbation Theory,” hep-th/0211249.
- [26] Y. Tachikawa, “Derivation of the Konishi anomaly relation from Dijkgraaf-Vafa with (bi-)fundamental matters,” hep-th/0211189.
- [27] S. G. Naculich, H. J. Schnitzer and N. Wyllard, “The $N = 2$ $U(N)$ gauge theory prepotential and periods from a perturbative matrix model calculation,” hep-th/0211123; “Matrix model approach to the $N=2$ $U(N)$

- gauge theory with matter in the fundamental representation”, hep-th/0211254
- [28] H. Itoyama and A. Morozov, “The Dijkgraaf-Vafa prepotential in the context of general Seiberg-Witten theory,” arXiv:hep-th/0211245.
 - [29] N. Seiberg, “Exact results on the space of vacua of four-dimensional SUSY gauge theories,” Phys. Rev. D **49**, 6857 (1994), hep-th/9402044.
 - [30] K. A. Intriligator, “Integrating in and exact superpotentials in 4-d,” Phys. Lett. B **336**, 409 (1994), hep-th/9407106].
 - [31] K. A. Intriligator and N. Seiberg, “Lectures on supersymmetric gauge theories and electric-magnetic duality,” Nucl. Phys. Proc. Suppl. **45BC**, 1 (1996), hep-th/9509066.
 - [32] D. Kutasov, “A Comment on duality in $N=1$ supersymmetric non-Abelian gauge theories,” Phys. Lett. B **351**, 230 (1995), hep-th/9503086; D. Kutasov and A. Schwimmer, hep-th/9505004; D. Kutasov, A. Schwimmer and N. Seiberg, Nucl. Phys. B **459**, 455 (1996), hep-th/9510222.
 - [33] C. Csaki and H. Murayama, “New confining $N = 1$ supersymmetric gauge theories,” Phys. Rev. D **59**, 065001 (1999), hep-th/9810014.
 - [34] P. C. Argyres, M. Ronen Plesser and N. Seiberg, “The Moduli Space of $N=2$ SUSY QCD and Duality in $N=1$ SUSY QCD,” Nucl. Phys. B **471**, 159 (1996), hep-th/9603042.
 - [35] K. Hori, H. Ooguri and Y. Oz, “Strong coupling dynamics of four-dimensional $N = 1$ gauge theories from M theory fivebrane,” Adv. Theor. Math. Phys. **1**, 1 (1998), hep-th/9706082.
 - [36] N. Seiberg, “Electric - magnetic duality in supersymmetric nonAbelian gauge theories,” Nucl. Phys. B **435**, 129 (1995), hep-th/9411149.
 - [37] C. Vafa, “Superstrings and topological strings at large N ,” J. Math. Phys. **42**, 2798 (2001), hep-th/0008142.
 - [38] M. Aganagic and C. Vafa, “Mirror symmetry and a $G(2)$ flop,” hep-th/0105225.

- [39] M. Aganagic and C. Vafa, “G(2) manifolds, mirror symmetry and geometric engineering,” hep-th/0110171.
- [40] M. Aganagic, M. Marino and C. Vafa, “All loop topological string amplitudes from Chern-Simons theory,” hep-th/0206164.
- [41] F. Cachazo, K. A. Intriligator and C. Vafa, “A large N duality via a geometric transition,” Nucl. Phys. B **603**, 3 (2001), hep-th/0103067.
- [42] J. D. Edelstein, K. Oh and R. Tatar, “Orientifold, geometric transition and large N duality for SO/Sp gauge theories,” JHEP **0105**, 009 (2001), hep-th/0104037.
- [43] F. Cachazo, S. Katz and C. Vafa, “Geometric transitions and N = 1 quiver theories,” hep-th/0108120.
- [44] F. Cachazo, B. Fiol, K. A. Intriligator, S. Katz and C. Vafa, “A geometric unification of dualities,” Nucl. Phys. B **628**, 3 (2002), hep-th/0110028.
- [45] B. Fiol, “Duality cascades and duality walls,” JHEP **0207**, 058 (2002), hep-th/0205155.
- [46] M. Aganagic, A. Klemm, M. Marino and C. Vafa, “Matrix model as a mirror of Chern-Simons theory,” arXiv:hep-th/0211098.
- [47] A. Klemm, M. Marino and S. Theisen, “Gravitational corrections in supersymmetric gauge theory and matrix models,” hep-th/0211216.
- [48] R. Dijkgraaf, A. Sinkovics and M. Temurhan, “Matrix Models and Gravitational Corrections,” hep-th/0211241.
- [49] K. Dasgupta, K. Oh and R. Tatar, “Geometric transition, large N dualities and MQCD dynamics,” Nucl. Phys. B **610**, 331 (2001), hep-th/0105066.
- [50] K. Dasgupta, K. Oh and R. Tatar, “Open/closed string dualities and Seiberg duality from geometric transitions in M-theory,” JHEP **0208**, 026 (2002), hep-th/0106040.

- [51] K. h. Oh and R. Tatar, “Duality and confinement in $N = 1$ supersymmetric theories from geometric transitions,” hep-th/0112040, to appear in Adv.Theor.Math.Phys.
- [52] K. Dasgupta, K. h. Oh, J. Park and R. Tatar, “Geometric transition versus cascading solution,” JHEP **0201**, 031 (2002), hep-th/0110050.
- [53] A. Hanany and E. Witten, “Type IIB superstrings, BPS monopoles, and three-dimensional gauge dynamics,” Nucl. Phys. B **492**, 152 (1997), hep-th/9611230.